

Display from CAplus

L# ANSWER 2 OF 4 CAPLUS COPYRIGHT 2003 ACS on STN

AN 1997:317775 CAPLUS

DN 126:295282

TI Fire-fighting aqueous emulsions consisting of water-insoluble fluorinated copolymers, ***surfactants***, fluorinated ***surfactants***, and water-miscible solvents

IN Garcia, Gilbert; Collette, Christian; Morillon, Elisabeth

PA Elf Atochem S.A., Fr.

SO Eur. Pat. Appl., 26 pp.

CODEN: EPXXDW

PATENT NO. KIND DATE APPLICATION NO. DATE

PI EP 765676 A1 19970402 EP 1996-401822 19960826

R: BE, CH, DE, DK, ES, FR, GB, IT, LI, NL, SE

FR 2739295 A1 19970404 FR 1995-11385 19950928

JP 09124884 A2 19970513 JP 1996-244969 19960917

AU 9665863 A1 19970410 AU 1996-65863 19960926

CA 2186773 AA 19970329 CA 1996-2186773 19960927

PRAI FR 1995-11385 A 19950928

AB Fire-fighting emulsions consist of aq. dispersions contg.: (1) 0.5-10 wt.% of a water-insol. fluorinated copolymer prep'd. from a monomer contg. a perfluorinated chain, a monomer contg. an ionic or ionizable chain, and/or a monomer contg. a nonionic chain, (2) 1-20 wt.% of a C5-18-branched or linear alkane ***surfactant***, (3) 0.5-10 wt.% of a fluorinated ***surfactant*** which, when present in aq. soln. at 1 g/L concn., gives rise to a surface tension at 20.degree. of 25 mN/m, and (4) 5-50 wt.% of a water-miscible solvent. The fluorinated copolymer is of general formula -[M1]x[M2]y[M3]z, in which M1 is a C5-20- ***perfluoroalkyl*** acrylate or methacrylate; M2 is an acrylic, methacrylic, or vinylic monomer with an ionizable or hydrophilic group; M3 is an acrylic, methacrylic, or vinylic monomer with a nonionic (esp. hydrophobic) group; and x = 50-95, yr = 1-25, and z = 0-10 (with an M1-M2 molar ratio of 1-10:1). A no. of ***surfactants*** are possible for the formulation, including ***betaines***, N,N-bis(2-carboxyethyl)amines, 2-alkyl(1-hydroxyethyl)imidazolines, quaternary ammonium chlorides, polyoxyalkylene monoethers, trialkylamine oxides, alkylbenzenesulfonic acids, etc.

IC ICM A62D001-00

DT Patent

LA French

EV979440213